

IMPACT OF PHARMACIST COUNSELING IN PATIENT KNOWLEDGE OF WARFARIN THERAPY

¹Dr. G. Sruthi, ²Dr. K. Manasa Reddy, ³Dr. K. Manasa

¹²³Assistant Professor

¹²³Department Of Pharmacy Practice

Vaagdevi Institute of Pharmaceutical Sciences, Bollikunta, Warangal. Telangana.

ABSTRACT

Background: The cornerstone of anticoagulant therapy is warfarin, although using it has been linked to a number of problems, most notably bleeding. Patients who are adequately informed about warfarin can reduce adverse medication responses and other related consequences while also improving anticoagulation control. The study's goal was to evaluate how patients' understanding of warfarin was impacted by their counselling from chemists. **Techniques:** A prospective randomised controlled trial was conducted at the National Heart Centre of Shahid Gangalal, Kathmandu, Nepal. Enrolling the individuals in the intervention and control groups was done using a straightforward random sample procedure.

The study involved 99 patients in all. To determine each group's baseline knowledge, the same questionnaire was given to them both. The intervention group received verbal counselling for twenty minutes, while the control group did not get any chemist intervention. After two months, a final evaluation of both groups' knowledge levels was conducted using the same questionnaire. **Findings:** Following the intervention, there was a statistically significant increase ($p < 0.05$) in the proportion of patients in the counselled group who knew enough about warfarin.

The findings also revealed that, following the intervention, a notably higher percentage of patients in the intervention group had improved INR (International Normalised Ratio) control. There was no discernible correlation between the patients' knowledge score, age, gender, or length of therapy. **Conclusion:** Patients' understanding of warfarin can be greatly enhanced by counselling from chemists.

Key words: Warfarin, INR, counselling, anticoagulation, and knowledge.

1. INTRODUCTION

Warfarin is the most widely recommended oral anticoagulant, used in the prevention of clotting among patients at risk for stroke or thromboembolic events.¹ The use of warfarin is challenged by its complex pharmacology and intrinsic risk of adverse outcomes, which requires routine INR (International Normalized ratio) monitoring to ensure safe and effective therapy.^{2,3} Clinicians have previously cited an unwillingness to prescribe warfarin due to its interactions with diet, drugs and need for monitoring,

all of which relies on patient compliance and treatment instructions and therefore is dependent on effective patient education.

Warfarin is a major contributor to potentially preventable drug-related adverse events, partly due to inadequate patient knowledge.⁵ A validated set of questionnaire provides a simple method of determining patients' knowledge of anticoagulation.⁶ Studies have concluded that patients' warfarin knowledge, a major contributor to anticoagulation control is poor and more attention should be given in educating

the patients' on warfarin use.⁷ Hence, it is of paramount importance for patients to have proper knowledge about the factors that may lead to warfarin related complications. Various studies have found that pharmacist provided verbal counseling improves warfarin knowledge.^{4,8,9} This can be accomplished by counseling so as to help prevent adverse health outcomes.⁴ However, studies assessing impact of educational intervention among Nepalese population is limited. Therefore, the study aims to assess the impact of pharmacist's counseling in

MATERIALS AND METHODS Study site and setting The study was conducted in the outpatient department of Shahid Gangalal National Heart Center, Kathmandu. This is a 200 bedded tertiary care hospital with an outpatient flow of 350 patients per day Study Duration The study was conducted for duration of 7 months from March to September 2015.

Selection of Participants Simple random sampling technique was used to assign the patients either to the intervention (pharmacist counseling) or the control group (usual care) where they had an equal chance of being allocated to either of the two groups. The patients selected in both the groups were similar in terms of socio-demographic variables. This ensured avoidance of patient selection bias in the study. The two homogenous groups were interviewed separately in different rooms and were kept unaware of their subjection to either intervention or control category to warrant achievement of blinding in the study. A patient who did not receive the pharmacist counseling but were similar to the intervention group in respect to other variables was the control group. All the patients who met the inclusion criteria and visited the outpatient pharmacy during the study duration

were enrolled in the study. A total of 99 patients (n=53, intervention group versus n=46, control group) were interviewed and 29 patients were lost to follow up.

Inclusion Criteria

- Patients visiting the outpatient pharmacy of Shahid Gangalal National Heart Center aged over 15 years.
- Patients who had been taking warfarin for at least 1 month.
- Patients willing to participate in the study.

Exclusion Criteria

- Patients who were recent users of warfarin (Less than 1 month).
- Patients admitted in the hospital.


2. Research Tool

The study was conducted using a questionnaire comprising of 15 questions on knowledge assessment, adapted from the pre-validated AKA (Anticoagulant Knowledge Assessment) Questionnaire and OAK (Oral Anticoagulant Knowledge) Questionnaire. The questionnaire was validated by pretesting. A pilot study was performed on 10 percent of the total expected sample size to strengthen the reliability and validity of the research. Each correct answer scored 1 point whereas an incorrect answer scored zero point. As a counseling aid, an information leaflet on warfarin therapy was provided to the patients in the intervention group. The information leaflet was compiled from "Oral Anticoagulant Therapy - Important information for patients" published by The National Patient Safety Agency, London and "The patient information, Warfarin Guide for patient and families" published by the John Hopkins Hospital. The information included in the leaflet was translated into Nepali language

which was checked and approved by an expert

Study Procedure A prospective randomized controlled study was conducted over a period of 7 months to assess the effect of pharmacist provided counseling in the knowledge of patients on warfarin. The study was approved by the ethical review board in Shahid Gangalal National Heart Center. The questionnaire was administered to the participants by the researcher himself. In the control group, the pharmacist did not intervene and only documented the baseline knowledge on warfarin therapy, on the first visit as well as on the subsequent visit after 2 months. In the intervention group the pharmacist obtained the baseline data and provided counseling to the patients for 20 min on warfarin therapy and its significance, dietary discipline, commonly observed adverse drug reactions and related preventive measures, significance of INR monitoring and warfarin drug interactions. A pharmacist prepared, leaflet containing necessary information on warfarin was also provided to the patients in the intervention group. After 2 months, the same questionnaire was used to assess the effect of counseling in the knowledge level of the patients who had received the educational intervention. In addition, INR (International Normalized Ratio) value in both the groups was collected at two time points, on the day of first interview and after 2 months. Twenty-nine patients were lost to follow up. Follow up data of only 36 patients in the intervention group and 34 patients in the control group could be obtained. Demographic details such as patients' age, gender, duration of warfarin therapy, INR value on the day of first interview and other clinically relevant information such as indication of warfarin therapy

was also collected in the patient performa.



A comparison of the number of patients with adequate knowledge in the intervention and the control group was done. Furthermore, a comparison of the number of patients with INR within the target goal was done, pre and post-intervention in the counseled group. Similarly, the difference in the number of patients with their INR value within the target range was also noted in the control group after 2 months. Disparity between percentages of patients answering each question correctly in the intervention and the control group was also observed. The data collected were entered in a separate spreadsheet of EXCEL and was analyzed using Statistical package for social sciences (spss) software version 20. (Figure 1) Statistical Analysis Application of the binomial distribution The questionnaire comprised of 15 questions with four options given as choice $p = \frac{1}{4}$ where, p is the probability of choosing the correct answer from the four options provided to the interviewee. If X is the random variable denoting number of correct answers provided by the interviewee, then X is distributed as a binomial with parameter $n = 15$ and $p = \frac{1}{4}$ Then $E(X)$

$= np = 15(1/4) = 3.75$ This implies that on average a person answers 4 questions correctly even if he/she has no knowledge of the subject. So, if $P(X \leq a) = 0.95$, the value of $a = 7$ was obtained from the binomial table provided $n=15$ and $p = 1/4$ so the $P(X \geq a) = 0.05$. This implies that if a person answers out of 15, 8 or more questions correctly we can conclude with 95% confidence that the person has adequate knowledge. Impact of pharmacist provided counseling in knowledge level of patients on warfarin was statistically analyzed using the Z test at 95% confidence interval. Similarly, separate evaluation of difference in number of patients with therapeutic INR after duration of 2 months in the intervention group as well as in the control group was done using the paired t-test. P value RESULTS Table 1 shows the demographic details of the study participants. A larger proportion of the patients were of the age group 31 to 50 years in both the intervention (40%, $n=21$) and the control group (39%, $n=18$). Counseled and the non-counseled group comprised of 64% ($n=34$) and 52% ($n=24$) of the female patients respectively. Most of the patients in our study had been taking warfarin for duration of more than 5 years. Table 2 illustrates the distribution of the study population as per the indication of warfarin. Majority of the male patients (32%, $n=13$) in our study had been taking warfarin for DVR (Double Valve Replacement) whereas the majority of the female patients had been taking warfarin for MVR (Mitral Valve Replacement) (36%, $n=21$). The most common indication for the use of anticoagulation was found to be MVR (30%, $n=30$).

Table 1: Characteristics of study population

Variables	Intervention Group (no. (%) of patients $n=53$)	Control Group (no. (%) of patients $n=46$)
Age (Years)		
15-30	10 (19)	16 (35)
31-50	21 (40)	18 (39)
51-65	17 (32)	10 (21)
65+	5 (9)	2 (4)
Gender		
Male	19 (36)	22 (48)
Female	34 (64)	24 (52)
Warfarin duration		
>6 months	11 (21)	8 (17)
6 months-1 year	4 (8)	3 (7)
1-5 years	18 (34)	13 (28)
>5 years	20 (38)	22 (48)

Table 2: Distribution of patients as per indication of warfarin

Indications	Male (no. (%) of patients) $n=41$	Female (no. (%) of patients) $n=58$
Aortic Valve Replacement	10 (22)	8 (14)
Double Valve Replacement	13 (32)	10 (17)
Deep Vein Thrombosis	3 (7)	6 (10)
Mitral Valve Replacement	9 (22)	21 (36)
Tricuspid Valve Replacement	6 (15)	13 (23)

Table 3: Evaluation of results of patients with adequate knowledge on warfarin in the intervention and control group

No. (%) of patients scoring more than 7 in follow up in control group ($n=34$)	No. (%) of patients scoring more than 7 after counseling in Interventional group ($n=36$)
13 (38)	32 (89)

3. RESULTS

Table 1 shows the demographic details of the study participants. A larger proportion of the patients were of the age group 31 to 50 years in both the intervention (40%, $n=21$) and the control group (39%, $n=18$). Counseled and the non-counseled group comprised of 64% ($n=34$) and 52% ($n=24$) of the female patients respectively. Most of the patients in our study had been taking warfarin for duration of more than 5 years. Table 2 illustrates the distribution of the study

population as per the indication of warfarin. Majority of the male patients (32%, n=13) in our study had been taking warfarin for DVR (Double Valve Replacement) whereas the majority of the female patients had been taking warfarin for MVR (Mitral Valve Replacement) (36%, n=21). The most common indication for the use of anticoagulation was found to be MVR (30%, n=30).

control group Table 3 shows a comparison of the number of patients with adequate knowledge on warfarin therapy in the intervention (38%, n=13) and the control group (89%, n=32). A significant improvement in warfarin knowledge was seen in the intervention group than in the control group ($p < 0.05$). Evaluation of results of patients with adequate knowledge on warfarin in intervention Group Table 4 shows the number of patients with adequate knowledge on warfarin therapy before counseling (22%, n=8), which rose to a figure of (89%, n=32) after counseling. A significant increment in the number was observed 2 months after counseling ($p < 0.05$). Table 5 illustrates the comparison of percentage of patients giving correct answers between the intervention and the control groups. It is representative of the data collected 2 months after the counseling. Majority of the questions were answered correctly in the counseled group when compared to the non-counseled group. Exception was the question related to interaction with alcohol (Question 5) and question related to interaction with vitamin (Question 9) where the non-counseled group achieved a slightly higher score than the counseled

Table 5. Percentage of patients giving correct answers in intervention and control group

Questions	Counseled No. (%) of patients (n=36)	Non-counseled No. (%) of patients (n=34)
Identifying different strengths of warfarin	18 (50)	17 (50)
Action in missing a dose	35 (97)	31 (91)
Food interaction	34 (94)	30 (88)
PT-INR means	35 (97)	28 (82)
Interaction with alcohol	12 (33)	12 (35)
Factor affecting warfarin action	18 (50)	48 (14)
Condition requiring emergency unit visit	27 (75)	15 (44)
Medicine of choice for headache relief	29 (80)	24 (70)
Interaction with vitamin	12 (33)	14 (41)
Fact about spinach intake	36 (100)	33 (97)
Most likely adverse effect of warfarin	67 (27)	2 (5)
When to take warfarin	36 (100)	34 (100)
Effect seen when INR is high	58 (16)	48 (14)
Effect necessitating to call Coumadin clinic	17 (47)	58 (17)
Length of time warfarin should be continued	36 (100)	34 (99)

group. All of the patients both in the counseled and the non-counseled group knew the best time of the day to take warfarin.

Table 6 displays the percentage of patients answering each question correctly before and after the counseling. The results showed an increase in the percentage of patients answering each question correctly post counseling. Furthermore, the results illustrated that a high percentage (70%, n=37) of the patients had less idea of identifying the different strengths of warfarin tablets (Question no.1). Knowledge concerning the administration time (Question no. 12) was also assessed. The proportion of patients who knew the correct administration time was almost (94%, n=50) even before counseling which rose to a perfect (100%, n=53) after pharmacist's counseling. Only (25%, n=13) of the patients knew about the action to be taken after missing a dose (Question no. 2) but the proportion increased to a whopping (97%, n=35) after counseling. An increase in mean knowledge score from 6.58 to 10.05 was observed after the intervention. The patient's knowledge was assessed on the following criteria: importance of INR monitoring, warfarin-food interactions, warfarin drugs interactions, adverse events of warfarin and actions to be taken when they occur and

Correlation between age, gender, duration of warfarin therapy and knowledge No significant association was found between age, gender, duration of warfarin therapy and the knowledge level of the patients ($p>0.05$) Assessment of the impact of pharmacist counseling on control of INR Comparison of the number of patients with INR within therapeutic range between the first day of interview and after 2 months, in the intervention and control groups is provided in Table 7 and Table 8 respectively. A significant increase in number of patients with INR within target range was obtained using paired t-test at 95% confidence interval in the intervention group ($n=30$, $p=0.005$) whereas no significant change was observed in the control group ($n=25$; $p=0.083$).

4. DISCUSSION

The study shows that majority of the warfarin users in both the intervention and control groups were of the

female patients (32% and 36%) respectively. (Table 2)

Pharmacist provided counseling can serve as a strong supportive measure to enhance warfarin related knowledge among the patients^{8,9} which further can serve as a major contributor to anticoagulation control.⁷ The results of this research also adheres to the above finding. This study examined whether pharmacist provided counseling improved the knowledge of patients regarding their warfarin therapy. Only (38%, $n=13$) of patients in the control group was found to have adequate knowledge on warfarin against a good number of (89%, $n=32$) in the intervention group. Table 3. Quite interestingly, a strikingly significant increase in percentage of patients with adequate knowledge was observed in the intervention group, 2 months post counseling (22%, $n=8$ vs. 89%; $n=32$). Table 4. This is consistent with the findings by Lakshmi et al. where mean knowledge score of patients' on anticoagulation rose significantly from 5.6($SD=3.2$) to 13.8($SD=0.94$) ($P=0.000$) after receiving counseling from clinical pharmacist, whereas no significant enhancement was observed in the control group ($p=0.218$).⁹ Another study by Collins et al. is reflective of similar findings where an increment of 34.9% in terms of number of patients achieving a passing score was noted after receiving verbal counseling on warfarin.⁸ But it is not in line with the result acquired by Hassan et al. which showed no significant differences in the knowledge of warfarin between the counseled and the non-counseled groups. However, the result by Hassan et al. is based on only 3 knowledge categories namely mechanism of action, interactions between alcohol and warfarin, side effects of warfarin. The reason for this difference probably is due to inclusion of 15 different knowledge assessment

Table 6. Percentage of patients correctly answering in intervention group

Questions	Pre intervention no (%) of patients (n=33)	Post intervention no(%) of Patients (n=36)
Identifying different strengths of warfarin	16 (30)	18 (50)
Action in missing a dose	13 (25)	35 (97)
Food interaction	16 (30)	34 (94)
PT-INR means	31 (58)	35 (97)
Interaction with alcohol	7 (13)	12 (33)
Factor affecting warfarin action	20 (38)	18 (50)
Condition requiring emergency unit visit	32 (61)	27 (75)
Medicine of choice for headache relief	28 (52)	29 (80)
Interaction with vitamin	7 (13)	12 (33)
Fact about spinach intake	47 (88)	36 (100)
Most likely adverse effect of warfarin	3 (5)	97 (27)
When to take warfarin	50 (94)	36 (100)
Effect seen when INR is high	4 (8)	58 (16)
Effect necessitating to call Coumadin clinic	8 (16)	17 (47)
Length of time warfarin should be continued	50 (94)	36 (100)

Table 7. INR control comparison in intervention group

No. (%of patients with INR in therapeutic range pre-counseling. (n=30)	No. of patients with INR in therapeutic range post-counseling. (n=30)
12(40)	24(80)

Table 8. INR control comparison in control group

No. (%of patients with INR in therapeutic range on first visit (n=25)	No. (%of patients with INR in therapeutic range on follow up (n=25)
12(48)	17(68)

age group 31-50 years (40% vs 39%). Percentage of female patients using warfarin was higher than their male counterparts in both the categories (64%, intervention group and 52%, control group). A larger proportion of patients had been under warfarin therapy for more than 5 years. (Table 1) The study revealed the most common indication for warfarin among male patients was DVR whereas it was MVR among the

criteria in the present study which might have possibly produced a result contrary to Hassan et al. 10 A positive impact of pharmacist provided counseling has also been demonstrated by the fact that most of the questions had been correctly addressed by patients in the intervention group as compared to those in the control group. (Table 5)

Number of patients with INR within therapeutic range rose significantly in the intervention group (n=30, p= 0.005) whereas no significant alteration was observed in the control group (n=25; p= 0.083). (Table 7, 8). This corroborates the findings of Khan et al. 11 who observed the percentage of time spent within the therapeutic range between two intervention groups and one control group. Although, no statistically significant difference was seen but the increase in percentage of time spent within the therapeutic range was relatively higher in the education plus self-monitoring group and the education alone group with a mean difference of 14.1 and 8.8 respectively, following 6 months of intervention. The findings of the present study backs the result of Khudair et al. 12 where significantly higher proportion of patients receiving warfarin education had controlled INR than the patients who did not receive any educational intervention (48%, n=116 vs. 26%, n=24; P=0.0136). Lakshmi et al. 9 has findings similar to ours where percentage of INRs within therapeutic range was found to be significantly greater in the intervention group than in the control group (77.4% vs. 46.5%; P=0.000). In regard to patients' comprehension of interaction of warfarin with food, initially (70% n=37) of patients were unaware of the foods that interfered with the working of warfarin. This number; however, reduced to (6%, n=2) two months after counseling. Approximate figures have

been presented in two of the previous studies.^{4,13} A great majority of patients (87%, n=46) believed in absence of interaction between alcohol and warfarin, which fortunately declined to (67%, n=24) after counseling. Yahaya et al. 14 and Shrestha et al. 13 reportedly presented a statistic of 34.6% and 44.1% respectively, of those who were ignorant of this potential interaction. Similarly, 48% of patients failed to select the safest non-steroidal anti-inflammatory drug while on warfarin, which after counseling lowered to 20%. A previous study reported a Figure of (44.1%, n=34) approximating to our data.¹³ Patients' ignorance on these essential factors may be ascribed to lack of adequate educational and assessment sessions in the hospitals. Nevertheless, proper counseling by pharmacists can alleviate the misconception and ignorance of patients regarding warfarin therapy.

No correlation was established between gender of the patients and warfarin knowledge. This is underpinned by the findings of previous studies.^{10,12-15} Similarly, warfarin knowledge was found to be unaffected by age. This is in contrary to the findings where age was found to be negatively correlated with total knowledge score of the patients.^{4,7,10,15,16} However, a study by Shrestha et al. 13 has findings similar to ours which found no association between age of the patients and warfarin knowledge. Furthermore, duration of therapy was also found to be uncorrelated with patient's knowledge which is similar to the results obtained by Baker et al. 17 and Hasan et al. 11 but opposes the findings of Shrestha S et al.

What this study adds to existing evidence? This study is first of its kind to address the impact of pharmacist provided counseling in knowledge of

warfarin as well as its aid in achieving INR control among the Nepalese patients. Findings of this study support the need and benefits of introducing the policies for physician pharmacist collaboration anticoagulation services in developing countries like Nepal.

Study Limitation Our study had a number of limitations. Firstly, this was a single institutional study and 29% of patients were lost to follow up. Secondly due to limited time frame we could provide only one session of counseling. Separate studies have suggested that no correlation exist between the patients' knowledge of warfarin therapy and anticoagulation control.^{15,18} However, the present study did not look into this matter. Although, the current study adds to the existing finding that educational intervention improves INR control, this result demands further investigation since only a single post INR value after 2 months of intervention was taken into account to assess the effect of counseling on INR. Additionally, INR values of only 30 patients in the intervention group and only 25 patients in the control group could be obtained at two time points.

Future Research Direction We realize that this subject merit further detailed exploration of impact of educational intervention and self-monitoring on INR control of patients on warfarin due to the aforementioned limitations.

5. CONCLUSION

Patients' who received pharmacists' counseling had better knowledge on warfarin therapy than those who did not receive the counseling. Our findings conclude that pharmacists' counseling aided by informational leaflets can play an important role in improving patients' warfarin related knowledge and probably have a remarkably positive

REFERENCES

1. Cabral KP, Ansell J, Hylek EM. Future directions of stroke prevention in atrial fibrillation: The potential impact of novel anticoagulants and stroke risk stratification. *J Thromb Haemost.* 2011;9(3):441-9. doi: 10.1111/j.1538-7836.2010.04179.x
2. Kim JH, Song YB, Shin DH, Kim JS, Choi JO, On YK, et al. How well does the target INR level maintains in warfarin-treated patients with non-valvular a trial Fibrillation. *Yonsei Med J.* 2009;50(1):83-8. doi:10.3349/ymj.2009.50.1.83
3. Bravata DM, Rosenbeck KR, Kancir S, Brass LM. The use of warfarin in veterans with atrial fibrillation. *BMC Cardiovasc Disord.* 2004;4(1):18. doi: 10.1186/1471-2261-4-18
4. Nasser S, Mullan J, Bajorek B. Challenges of older patients' knowledge about warfare therapy. *J Prim Care Community Health.* 2012;3(1):56-74. doi: 10.1177/2150131911416365
5. Cheah GM, Martens KH. Coumadin knowledge deficits: Do recently hospitalized patients know how to safely manage the medication?. *Home Healthc Nurse.* 2003;21(2):94-100.
6. Taylor FC, Ramsey ME, Tan G, Gabbay J, Cohen H. Evaluation of patients' knowledge about anticoagulation treatment. *Qual Health Care.* 1994;3(2):79-85.
7. Tang EO, Lai CS, Lee KK, Wong RS, Cheng G, Chan TY. Relationship between patients' warfarin knowledge and anticoagulation control. *Ann Pharmacother.* 2003;37(1):34-9.
8. Collins S, Barber A, Sahm LJ. Pharmacist's counselling improves patient knowledge regarding warfarin, irrespective of health literacy level. *Pharmacy.* 2014;2(1):114-23. doi: 10.3390/pharmacy2010114
9. Lakshmi R, James E, Kirthivasan R. Study on

- impact of clinical pharmacist's interventions in the optimal use of oral anticoagulants in stroke patients. *Indian J Pharm Sci.* 2013;75(1):53-9. doi:10.4103/0250-474X.113550 10. Hasan SS, Shamala R, Syed IA, Basariah N, Chong DW, Mei TK, et al. Factors affecting warfarin-related knowledge and INR control of patients attending physician- and pharmacist-managed anticoagulation clinics. *J Pharm Pract.* 2011;24(5):485-93. doi: 10.1177/0897190011415684 11. Khan TI, Kamali F, Kesteven P, Avery P, Wynne H. The value of education and self-monitoring in the management of warfarin therapy in older patients with unstable control of anticoagulation. *Br J Haematol.* 2004;126(4):557-64. doi:10.1111/j.13652141.2004.05074.x 12. Khudair IF, Hanssens YI. Evaluation of patients' knowledge on warfarin in outpatient anticoagulation clinics in a teaching hospital in Qatar. *Saudi Med J.* 2010;31(6):672-7. 13. Shrestha S, Sapkota B, Kumpakha A, Acharya U, Sharma R. Evaluation of patients' knowledge on warfarin in outpatient pharmacy of a tertiary care cardiac center. *BMC Res Notes.* 2015;8(1):429. doi: 10.1186/s13104-015-1416-1 14. Yahaya AH, Hassali MA, Awaisu A, Shafie AA. Factors associated with warfarin therapy knowledge and anticoagulation control among patients attending a warfarin clinic in Malaysia. *JCDR.* 2009;3(4):1663-70. 15. Hu A, Chow CM, Dao D, Errett L, Keith M. Factors influencing patient knowledge of warfarin therapy after mechanical heart valve replacement. *J Cardiovasc Nurs.* 2006;21(3):169-75. 16. Dantas GC, Thompson BV, Manson JA, Tracy CS, Upshur RE. Patients' perspectives on taking warfarin: Qualitative study in family practice. *BMC Fam Pract.* 2004;5(1):15. doi: 10.1186/1471-2296-5-15. 17. Baker JW, Pierce KL, Ryals CA. INR Goal attainment and oral anticoagulation knowledge of patients enrolled in an anticoagulation clinic in a Veterans Affairs medical center. *J Manag Care Pharm.* 2011;17(2):133-42. doi: 10.18553/jmcp.2011.17.2.13 18. Davis NJ, Billett HH, Cohen HW, Arnsten JH. Impact of adherence, knowledge and quality of life on anticoagulation control. *Ann Pharmacother.* 2005;39(4):632-6.